

Briefing 1621

Ash dieback disease

Summary

Ash dieback is a disease of ash trees caused by a fungus called *Chalara fraxinea*. The disease causes leaf loss and crown dieback in affected trees and it may lead to tree death. Ash trees suffering from symptoms likely to be caused by *Chalara fraxinea* are increasingly being found across Europe. *Chalara fraxinea* is being treated as a quarantine pest under national emergency measures; it is important that suspected cases of the disease are reported

This paper is summarised from a Forestry Commission Pest Alert. 'Ash dieback disease'. The original paper has helpful illustrations and can be accessed at:

[http://www.forestry.gov.uk/pdf/pest-alert-ash-dieback-2012.pdf/\\$FILE/pest-alert-ash-dieback-2012.pdf](http://www.forestry.gov.uk/pdf/pest-alert-ash-dieback-2012.pdf/$FILE/pest-alert-ash-dieback-2012.pdf)

A number of countries in Europe have reported trees with ash dieback disease. Common ash (*Fraxinus excelsior*) is the most frequently affected species although *Fraxinus angustifolia* and the 'Pendula' ornamental variety of common ash have also been reported as hosts. Susceptibility of other ash species is unknown. Ash trees of all ages can be affected, but it has been reported that mortality is particularly common in saplings.

Symptoms to look out for

Symptoms of *Chalara fraxinea* can be visible on leaves, shoots and branches of affected trees. In severe cases, the entire crown shows leaf loss and dieback and there may also be the formation of epicormic shoots on branches and the trunk.

Leaves can suffer from wilting and black-brownish discoloration at the leaf base and midrib. Dieback of shoots and twigs is also very characteristic.

Small lens-shaped lesions or necrotic spots appear on the bark of stems and branches and enlarge to form perennial cankers. These cause wilting and dieback of shoots and branches, particularly in the upper crown. Underneath the bark lesions, the wood has a brownish to grey discoloration which often extends longitudinally beyond the bark necrosis.

Trees with withered tops and shoots are very characteristic. Heavily affected trees have extensive shoot, twig and branch dieback and often have prolific epicormic shoots. *Chalara fraxinea* has also been isolated from the roots of symptomatic trees, as well as from leaves, shoots and branch/stem lesions.

How the disease spreads

Details of how *Chalara fraxinea* is spread are uncertain, but local spread may be via rain splash or even transmission by insects. Over longer distances the risk of disease spread is most likely to be via the

movement of diseased ash plants. Movement of logs or unsawn wood from infected trees may also be a pathway for the disease.

Other disorders of ash trees

There are a number of disorders of ash trees in Britain that may be mistaken for infection by *Chalara fraxinea*:

- A disorder also known as 'ash dieback' which has been known for many years in Britain and is widespread. Affected trees typically suffer from severe crown dieback but there may be recovery growth in the form of secondary shoots and 'epicormics'. The cause is thought to be mainly physical rather than biological. Large, mature trees are most often affected, and crown symptoms are most marked in trees adjacent to arable land where root disturbance is thought to play a part in the decline.
- Cankers on ash stems can be caused by the common fungal pathogen *Nectria galligena*, and the ash bark beetle *Leperisinus varius* can also cause bark necrosis.
- The activities of the ash bud moth *Prays fraxinella* in spring can also be mistaken for infection by *Chalara fraxinea* as the moth larvae mine into the base of shoots causing them to wilt and die.
- Frost and drought can cause similar damage in the form of dead shoots on ash trees.

Who to contact

Please report suspected cases of the disease to:

- Forest Research Disease Diagnostic Advisory Service T: 01420 23000; E: ddas.ah@forestry.gsi.gov.uk
- Forestry Commission Plant Health Service T: 0131 314 6414; E: plant.health@forestry.gsi.gov.uk
- Fera Plant Health and Seeds Inspectorate T: 01904 465625; E: planthealth.info@fera.gsi.gov.uk

Alan Spedding, 29 October 2012

RuSource briefings provide concise information on current farming and rural issues for rural professionals. They are circulated weekly by email and produced by Alan Spedding in association with the Arthur Rank Centre, the national focus for the rural church. Previous briefings can be accessed on the Arthur Rank Centre website at <http://www.arthurrankcentre.org.uk/publications-and-resources/rusource>

RuSource is a voluntary project partly supported by donations and sponsorship.

© Alan Spedding 2012. This briefing may be reproduced or transmitted in its entirety free of charge. Where extracts are used, their source must be acknowledged. RuSource briefings may not be reproduced in any publication or offered for sale without the prior permission of the copyright holder.

If you would like to be put on the list for regular briefings or have any other queries about the service contact alan.spedding@btopenworld.com.